



Contents lists available at SciVerse ScienceDirect

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laa

A simple proof of the generalized Craig–Sakamoto theorem

Jin Zhang^{a,*}, Jikun Yi^b^a School of Mathematics and Statistics, Yunnan University, Kunming, Yunnan 650091, China^b College of Continued Education, Yunnan University, Kunming, Yunnan 650091, China

ARTICLE INFO

Article history:

Received 20 September 2011

Accepted 26 February 2012

Available online 16 April 2012

Submitted by H. Schneider

AMS classification:

15A15

15A18

15A24

15A48

Keywords:

Eigenvalues

Independence

Quadratic forms

Symmetric matrices

ABSTRACT

The Craig–Sakamoto theorem establishes the independence of two quadratic forms in normal variates. In this article, we provide a simple proof of a generalized Craig–Sakamoto theorem.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

The Craig–Sakamoto theorem establishes the independence of two quadratic forms in normal random variables. For its history, refer to [3,7,8]. After some development, the Craig–Sakamoto theorem can be stated as the following mathematical form.

Theorem 1 (Craig–Sakamoto). *Let A and B be $n \times n$ real symmetric matrices. Then $|I_n - xA - yB| = |I_n - xA||I_n - yB|$ for any $x, y \in \mathbf{R}$ if and only if $AB = 0$.*

Many proofs can be found in [1,2,4–6,9]. In the next section, we provide a simple proof of a generalized Craig–Sakamoto theorem below.

* Corresponding author.

E-mail addresses: jinzhang@ynu.edu.cn (J. Zhang), jikunyi@netscape.net (J. Yi).

Theorem 2. Let A and B be $n \times n$ real symmetric matrices with a_1, \dots, a_s and b_1, \dots, b_t as their nonzero eigenvalues. Then the following conditions are equivalent.

- (1) $AB = 0$.
- (2) $|I_n - xA - yB| = |I_n - xA||I_n - yB|$ for any $x, y \in \mathbf{R}$.
- (3) $|I_n - x(A + B)| = |I_n - xA||I_n - xB|$ for any $x \in \mathbf{R}$.
- (4) The nonzero eigenvalues of $A + B$ are $\{a_1, \dots, a_s, b_1, \dots, b_t\}$.

2. The proof

The proofs for (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) are straightforward. To prove (4) \Rightarrow (1), note that the rank $r(A + B) = s + t$ and $|U + V| \geq |U| + |V|$ if U and V are positive semi-definite.

Case (a) $s + t = n$: Let $A_1 = \text{diag}(a_1, \dots, a_s)$ and $B_1 = \text{diag}(b_1, \dots, b_t)$. Without loss of generality, assume that $A = \begin{pmatrix} A_1 & 0 \\ 0 & 0 \end{pmatrix}$, $B = T \begin{pmatrix} 0 & 0 \\ 0 & B_1 \end{pmatrix} T'$ and $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix}$ is orthogonal. Then $A + B = \begin{pmatrix} I & T_2 \\ 0 & T_4 \end{pmatrix} \begin{pmatrix} A_1 & 0 \\ 0 & B_1 \end{pmatrix} \begin{pmatrix} I & 0 \\ T_2' & T_4' \end{pmatrix}$, $|A + B| = |A_1||B_1||T_4'T_4|$ and $|T_4'T_4| = 1 = |I_t| = |T_2'T_2 + T_4'T_4| \geq |T_2'T_2| + |T_4'T_4|$. Thus $T_2 = 0$ and $AB = 0$.

Case (b) $s + t < n$: $r \begin{pmatrix} A \\ B \end{pmatrix} = s + t$ since $r(A + B) \leq r \begin{pmatrix} A \\ B \end{pmatrix} \leq r(A) + r(B)$.

Let $P = (x_1, x_2, \dots, x_n)$ be an orthogonal matrix, where $\begin{pmatrix} A \\ B \end{pmatrix} x_i = 0$, $i = 1, 2, \dots, n - s - t$. Then, $P'AP = \begin{pmatrix} 0 & 0 \\ 0 & A^* \end{pmatrix}$, $P'BP = \begin{pmatrix} 0 & 0 \\ 0 & B^* \end{pmatrix}$ and $P'(A + B)P = \begin{pmatrix} 0 & 0 \\ 0 & A^* + B^* \end{pmatrix}$, where A^* and B^* satisfy (4) and Case (a). Thus $A^*B^* = 0$ and $AB = 0$.

Acknowledgements

This research is supported by the Natural Science Foundation of China (NSFC, Grant No. 1161054). The author thank a referee for helpful comments and suggestions.

References

- [1] H. Carriau, Close to the Craig–Sakamoto theorem, *Linear Algebra Appl.* 432 (2010) 777–779.
- [2] H. Carriau, P. Lassère, One more simple proof of the Craig–Sakamoto theorem, *Linear Algebra Appl.* 431 (2009) 1616–1619.
- [3] M.F. Driscoll, W.R. Gundberg Jr., A history of the development of Craig's theorem, *Amer. Statist.* 42 (1986) 139–142.
- [4] C.-K. Li, A simple proof of the Craig–Sakamoto theorem, *Linear Algebra Appl.* 321 (2000) 281–283.
- [5] M. Marcus, On a determinant result of I. Olkin, *Linear Algebra Appl.* 277 (1998) 237–238.
- [6] M. Matsuura, On the Craig–Sakamoto theorem and Olkin's determinantal result, *Linear Algebra Appl.* 364 (2003) 321–323.
- [7] J. Ogawa, A history of the development of Craig–Sakamoto's theorem viewed from the Japanese standpoint, *Proc. Ann. Inst. Statist. Math.* 41 (1993) 47–59.
- [8] J. Ogawa, I. Olkin, A tale of two countries: the Craig–Sakamoto–Matusita theorem, *J. Statist. Plann. Infer.* 138 (2008) 3419–3428.
- [9] I. Olkin, A determinantal proof of the Craig–Sakamoto theorem, *Linear Algebra Appl.* 264 (1997) 217–223.